

TITLE OF THE INVENTION

NOVEL AGGRECANASE MOLECULES

The present invention relates to the discovery of nucleotide sequences encoding novel aggrecanase molecules, the aggrecanase proteins and processes for producing them. The invention further relates to the development of inhibitors of, as well as antibodies to the aggrecanase enzymes. These inhibitors and antibodies may be useful for the treatment of various aggrecanase-associated conditions including osteoarthritis.

BACKGROUND OF THE INVENTION

Aggrecan is a major extracellular component of articular cartilage. It is a proteoglycan responsible for providing cartilage with its mechanical properties of compressibility and elasticity. The loss of aggrecan has been implicated in the degradation of articular cartilage in arthritic diseases. Osteoarthritis is a debilitating disease which affects at least 30 million Americans [MacLean et al. *J Rheumatol* 25:2213-8. (1998)]. Osteoarthritis can severely reduce quality of life due to degradation of articular cartilage and the resulting chronic pain. An early and important characteristic of the osteoarthritic process is loss of aggrecan from the extracellular matrix [Brandt, KD. and Mankin HJ. *Pathogenesis of Osteoarthritis*, in Textbook of Rheumatology, WB Saunders Company, Philadelphia, PA pgs. 1355-1373. (1993)]. The large, sugar-containing portion of aggrecan is thereby lost from the extra-cellular matrix, resulting in deficiencies in the biomechanical characteristics of the cartilage.

A proteolytic activity termed "aggrecanase" is thought to be responsible for the cleavage of aggrecan thereby having a role in cartilage degradation associated with osteoarthritis and inflammatory joint disease. Work has been conducted to identify the enzyme responsible for the degradation of aggrecan in human osteoarthritic cartilage.

- 5 Two enzymatic cleavage sites have been identified within the interglobular domain of aggrecan. One (Asn³⁴¹-Phe³⁴²) is observed to be cleaved by several known metalloproteases [Flannery, CR et al. J Biol Chem 267:1008-14. 1992; Fosang, AJ et al. Biochemical J. 304:347-351. (1994)]. The aggrecan fragment found in human synovial fluid, and generated by IL-1 induced cartilage aggrecan cleavage is at the Glu³⁷³-Ala³⁷⁴
- 10 bond [Sandy, JD, et al. J Clin Invest 69:1512-1516. (1992); Lohmander LS, et al. Arthritis Rheum 36: 1214-1222. (1993); Sandy JD et al. J Biol Chem. 266: 8683-8685. (1991)], indicating that none of the known enzymes are responsible for aggrecan cleavage in vivo.

- Recently, identification of two enzymes, aggrecanase-1(ADAMTS 4) and
- 15 aggrecanase -2 (ADAMTS-11) within the "Disintegrin-like and Metalloprotease with Thrombospondin type 1 motif" (ADAM-TS) family have been identified which are synthesized by IL-1 stimulated cartilage and cleave aggrecan at the appropriate site [Tortorella MD, et al. Science 284:1664-6. (1999); Abbaszade, I, et al. J Biol Chem 274: 23443-23450. (1999)]. It is possible that these enzymes could be synthesized by
- 20 osteoarthritic human articular cartilage. It is also contemplated that there are other, related enzymes in the ADAM-TS family which are capable of cleaving aggrecan at the Glu³⁷³-Ala³⁷⁴ bond and could contribute to aggrecan cleavage in osteoarthritis.

SUMMARY OF THE INVENTION

The present invention is directed to the identification of aggrecanase protein molecules capable of cleaving aggrecanase, the nucleotide sequences which encode the aggrecanase enzymes, and processes for the production of aggrecanases. These enzymes
5 are contemplated to be characterized as having proteolytic aggrecanase activity. The invention further includes compositions comprising these enzymes as well as antibodies to these enzymes. In addition, the invention includes methods for developing inhibitors of aggrecanase which block the enzyme's proteolytic activity. These inhibitors and antibodies may be used in various assays and therapies for treatment of conditions
10 characterized by the degradation of articular cartilage.

The nucleotide sequence of the aggrecanase molecule of the present invention is set forth in SEQ ID NO: 3. In another embodiment, the nucleotide sequence of the aggrecanase molecule of the present invention is set forth SEQ ID NO: 1 from nucleotide
1 to #3766. In another embodiment the nucleotide sequence of the invention
15 comprises nucleotide #1086(TCG) to # 3396 (CGC) of SEQ ID NO: 1. The invention further includes equivalent degenerative codon sequences of the sequences set forth in SEQ ID NO: 1, as well as fragments thereof which exhibit aggrecanase activity.

The amino acid sequence of an isolated aggrecanase molecule of the invention comprises the sequence set forth in SEQ. ID. NO:4. The amino acid sequence of an
20 isolated aggrecanase molecule comprises the sequence set forth in SEQ ID. No. 2. The invention further includes fragments of the amino acid sequence which encode molecules exhibiting aggrecanase activity.

The human aggrecanase protein or a fragment thereof may be produced by culturing a cell transformed with a DNA sequence of SEQ ID NO: 3 or SEQ ID NO: 1 comprising nucleotide # 1 to #3766 of SEQ ID NO: 1 or comprising nucleotide # 1086 to #3396 of SEQ ID NO:1 and recovering and purifying from the culture medium a protein
5 characterized by the amino acid sequence set forth in SEQ ID NO: 4 or SEQ ID NO: 2, respectively, substantially free from other proteinaceous materials with which it is co-produced. For production in mammalian cells, the DNA sequence further comprises a DNA sequence encoding a suitable propeptide 5' to and linked in frame to the nucleotide sequence encoding the aggrecanase enzyme.

10 The invention includes methods for obtaining additional aggrecanase molecules, the DNA sequence obtained by this method and the protein encoded thereby. The method for isolation of the full length sequence involves utilizing the aggrecanase sequence set forth in SEQ ID NO: 3 or SEQ ID NO: 1 from nucleotide # 1086 to #3396 to design probes for screening using standard procedures known to those skilled in the art.

15 It is expected that other species have DNA sequences homologous to human aggrecanase enzyme. The invention, therefore, includes methods for obtaining the DNA sequences encoding other aggrecanase molecules, the DNA sequences obtained by those methods, and the protein encoded by those DNA sequences. This method entails utilizing the nucleotide sequence of the invention or portions thereof to design probes to screen libraries for the
20 corresponding gene from other species or coding sequences or fragments thereof from using standard techniques. Thus, the present invention may include DNA sequences from other species, which are homologous to the human aggrecanase protein and can be obtained using the human sequence. The present invention may also include functional fragments of the aggrecanase

protein, and DNA sequences encoding such functional fragments, as well as functional fragments of other related proteins. The ability of such a fragment to function is determinable by assay of the protein in the biological assays described for the assay of the aggrecanase protein.

The aggrecanase proteins of the present invention may be produced by culturing
5 a cell transformed with the DNA sequence of SEQ ID NO: 3 or SEQ ID NO: 1
comprising nucleotide # 1 to # 3766 of SEQ ID NO: 1 or comprising nucleotide # 1086
to #3396 of SEQ ID NO 1 and recovering and purifying aggrecanase protein from the
culture medium. In one embodiment the protein comprises amino acid sequence of SEQ
ID NO: 4 or amino acid #1 to #770 of SEQ ID No:2. The purified expressed protein is
10 substantially free from other proteinaceous materials with which it is co-produced, as well
as from other contaminants. The recovered purified protein is contemplated to exhibit
proteolytic aggrecanase activity cleaving aggrecan. Thus, the proteins of the invention
may be further characterized by the ability to demonstrate aggrecan proteolytic activity in
an assay which determines the presence of an aggrecan-degrading molecule. These
15 assays or the development thereof is within the knowledge of one skilled in the art. Such
assays may involve contacting an aggrecan substrate with the aggrecanase molecule and
monitoring the production of aggrecan fragments [see for example, Hughes et al.,
Biochem J 305: 799-804(1995); Mercuri et al, J. Bio Chem. 274:32387-32395 (1999)]

In another embodiment, the invention includes methods for developing inhibitors
20 of aggrecanase and the inhibitors produced thereby. These inhibitors prevent cleavage of
aggrecan. The method may entail the determination of binding sites based on the three
dimensional structure of aggrecanase and aggrecan and developing a molecule reactive
with the binding site. Candidate molecules are assayed for inhibitory activity. Additional

standard methods for developing inhibitors of the aggrecanase molecule are known to those skilled in the art. Assays for the inhibitors involve contacting a mixture of aggrecan and the inhibitor with an aggrecanase molecule followed by measurement of the aggrecanase inhibition, for instance by detection and measurement of aggrecan fragments
5 produced by cleavage at an aggrecanase susceptible site.

Another aspect of the invention therefore provides pharmaceutical compositions containing a therapeutically effective amount of aggrecanase inhibitors, in a pharmaceutically acceptable vehicle.

Aggrecanase-mediated degradation of aggrecan in cartilage has been implicated in
10 osteoarthritis and other inflammatory diseases. Therefore, these compositions of the invention may be used in the treatment of diseases characterized by the degradation of aggrecan and/or an upregulation of aggrecanase. The compositions may be used in the treatment of these conditions or in the prevention thereof.

The invention includes methods for treating patients suffering from conditions
15 characterized by a degradation of aggrecan or preventing such conditions. These methods, according to the invention, entail administering to a patient needing such treatment, an effective amount of a composition comprising an aggrecanase inhibitor which inhibits the proteolytic activity of aggrecanase enzymes.

Still a further aspect of the invention are DNA sequences coding for expression of
20 an aggrecanase protein. Such sequences include the sequence of nucleotides in a 5' to 3' direction illustrated in SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO: 1 or SEQ ID NO:3 and DNA sequences

which, but for the degeneracy of the genetic code, are identical to the DNA sequence of SEQ ID NO: 1 and SEQ ID NO: 3, and encode an aggrecanase protein. Further included in the present invention are DNA sequences which hybridize under stringent conditions with the DNA sequence of SEQ ID NO: 1 and SEQ ID NO: 3 and encode a protein having the ability to cleave aggrecan. Preferred DNA sequences include those which hybridize under stringent conditions [see, T. Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389]. It is generally preferred that such DNA sequences encode a polypeptide which is at least about 80% homologous, and more preferably at least about 90% homologous, to the sequence of set forth in SEQ ID NO: 3 or SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO: 1. Finally, allelic or other variations of the sequences of SEQ ID NO: 1 or SEQ ID NO: 3, whether such nucleotide changes result in changes in the peptide sequence or not, but where the peptide sequence still has aggrecanase activity, are also included in the present invention. The present invention also includes fragments of the DNA sequence shown in SEQ ID NO: 1 which encode a polypeptide which retains the activity of aggrecanase.

The DNA sequences of the present invention are useful, for example, as probes for the detection of mRNA encoding aggrecanase in a given cell population. Thus, the present invention includes methods of detecting or diagnosing genetic disorders involving the aggrecanase, or disorders involving cellular, organ or tissue disorders in which aggrecanase is irregularly transcribed or expressed. The DNA sequences may also be useful for preparing vectors for gene therapy applications as described below.

A further aspect of the invention includes vectors comprising a DNA sequence as described above in operative association with an expression control sequence therefor. These vectors may be employed in a novel process for producing an aggrecanase protein of the invention in which a cell line transformed with a DNA sequence encoding an

5 aggrecanase protein in operative association with an expression control sequence therefor, is cultured in a suitable culture medium and an aggrecanase protein is recovered and purified therefrom. This process may employ a number of known cells both prokaryotic and eukaryotic as host cells for expression of the polypeptide. The vectors may be used in gene therapy applications. In such use, the vectors may be transfected into

10 the cells of a patient *ex vivo*, and the cells may be reintroduced into a patient. Alternatively, the vectors may be introduced into a patient *in vivo* through targeted transfection.

Still a further aspect of the invention are aggrecanase proteins or polypeptides. Such polypeptides are characterized by having an amino acid sequence including the

15 sequence illustrated in SEQ ID NO. 2 or 4, variants of the amino acid sequence of SEQ ID NO.2 or 4, including naturally occurring allelic variants, and other variants in which the protein retains the ability to cleave aggrecan characteristic of aggrecanase molecules. Preferred polypeptides include a polypeptide which is at least about 80% homologous, and more preferably at least about 90% homologous, to the amino acid sequence shown in

20 SEQ ID NO. 2 or 4. Finally, allelic or other variations of the sequences of SEQ ID NO. 2 or 4, whether such amino acid changes are induced by mutagenesis, chemical alteration, or by alteration of DNA sequence used to produce the polypeptide, where the peptide

sequence still has aggrecanase activity, are also included in the present invention. The present invention also includes fragments of the amino acid sequence of SEQ ID NO. 2 or 4 which retain the activity of aggrecanase protein.

The purified proteins of the present inventions may be used to generate antibodies, either monoclonal or polyclonal, to aggrecanase and/or other aggrecanase-related proteins, using methods that are known in the art of antibody production. Thus, the present invention also includes antibodies to aggrecanase or other related proteins. The antibodies may be useful for detection and/or purification of aggrecanase or related proteins, or for inhibiting or preventing the effects of aggrecanase. The aggrecanase of the invention or portions thereof may be utilized to prepare antibodies that specifically bind to aggrecanase.

DETAILED DESCRIPTION OF THE INVENTION

The human aggrecanase of the present invention comprises the nucleotide sequence set in SEQ ID NO: 3. In another embodiment, the human aggrecanase of the present invention comprises nucleotides # 1 to # 3766 or nucleotides # 1086 to #3396 of SEQ ID NO 1. The human aggrecanase protein sequence comprises the amino acid sequence set forth in SEQ ID NO: 4. In another embodiment, the human aggrecanase protein sequence comprises amino acids # 1 to # 770 set forth in SEQ ID NO. 2. Further sequences of the aggrecanase of the present invention may be obtained using the sequences of SEQ ID NO: 3 or SEQ ID NO. 1 comprising nucleotides # 1086 to # 3396 to design probes for screening for the full sequence using standard techniques.

The aggrecanase proteins of the present invention, include polypeptides comprising the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4 and having the ability to cleave aggrecan.

The aggrecanase proteins recovered from the culture medium are purified by

5 isolating them from other proteinaceous materials from which they are co-produced and from other contaminants present. The isolated and purified proteins may be characterized by the ability to cleave aggrecan substrate. The aggrecanase proteins provided herein also include factors encoded by the sequences similar to those of SEQ ID NO: 3 or the sequence of SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or

10 comprising nucleotide # 1086 to #3396 of SEQ ID NO 1, but into which modifications or deletions are naturally provided (e.g. allelic variations in the nucleotide sequence which may result in amino acid changes in the polypeptide) or deliberately engineered. For example, synthetic polypeptides may wholly or partially duplicate continuous sequences of the amino acid residues of SEQ ID NO. 2 or SEQ ID NO: 4. These sequences, by

15 virtue of sharing primary, secondary, or tertiary structural and conformational characteristics with aggrecanase molecules may possess biological properties in common therewith. It is known, for example that numerous conservative amino acid substitutions are possible without significantly modifying the structure and conformation of a protein, thus maintaining the biological properties as well. For example, it is recognized that

20 conservative amino acid substitutions may be made among amino acids with basic side chains, such as lysine (Lys or K), arginine (Arg or R) and histidine (His or H), amino acids with acidic side chains, such as aspartic acid (Asp or D) and glutamic acid (Glu or

E); amino acids with uncharged polar side chains, such as asparagine (Asn or N), glutamine (Gln or Q), serine (Ser or S), threonine (Thr or T), and tyrosine (Tyr or Y); and amino acids with nonpolar side chains, such as alanine (Ala or A), glycine (Gly or G), valine (Val or V), leucine (Leu or L), isoleucine (Ile or I), proline (Pro or P),

5 phenylalanine (Phe or F), methionine (Met or M), tryptophan (Trp or W) and cysteine (Cys or C). Thus, these modifications and deletions of the native aggrecanase may be employed as biologically active substitutes for naturally-occurring aggrecanase and in the development of inhibitors other polypeptides in therapeutic processes. It can be readily determined whether a given variant of aggrecanase maintains the biological activity of

10 aggrecanase by subjecting both aggrecanase and the variant of aggrecanase, as well as inhibitors thereof, to the assays described in the examples.

Other specific mutations of the sequences of aggrecanase proteins described herein involve modifications of glycosylation sites. These modifications may involve O-linked or N-linked glycosylation sites. For instance, the absence of glycosylation or only

15 partial glycosylation results from amino acid substitution or deletion at asparagine-linked glycosylation recognition sites. The asparagine-linked glycosylation recognition sites comprise tripeptide sequences which are specifically recognized by appropriate cellular glycosylation enzymes. These tripeptide sequences are either asparagine-X-threonine or asparagine-X-serine, where X is usually any amino acid. A variety of amino acid

20 substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) results in non-glycosylation at the modified tripeptide sequence. Additionally, bacterial

expression of aggrecanase-related protein will also result in production of a non-glycosylated protein, even if the glycosylation sites are left unmodified.

The present invention also encompasses the novel DNA sequences, free of association with DNA sequences encoding other proteinaceous materials, and coding for expression of aggrecanase proteins. These DNA sequences include those depicted in SEQ ID NO: 1 or 3 in a 5' to 3' direction and those sequences which hybridize thereto under stringent hybridization washing conditions [for example, 0.1X SSC, 0.1% SDS at 65°C; see, T. Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389] and encode a protein having aggrecanase proteolytic activity. These DNA sequences also include those which comprise the DNA sequence of SEQ ID NO: 1 and those which hybridize thereto under stringent hybridization conditions and encode a protein which maintain the other activities disclosed for aggrecanase.

Similarly, DNA sequences which code for aggrecanase proteins coded for by the sequences of SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO 1 or the sequence of SEQ ID NO: 3 or aggrecanase proteins which comprise the amino acid sequence of SEQ ID NO. 2 or 4, but which differ in codon sequence due to the degeneracies of the genetic code or allelic variations (naturally-occurring base changes in the species population which may or may not result in an amino acid change) also encode the novel factors described herein.

Variations in the DNA sequences of SEQ ID NO: 3 or SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO 1 which are caused by point mutations or by induced modifications (including insertion,

deletion, and substitution) to enhance the activity, half-life or production of the polypeptides encoded are also encompassed in the invention.

Another aspect of the present invention provides a novel method for producing aggrecanase proteins. The method of the present invention involves culturing a suitable cell line, which has been transformed with a DNA sequence encoding a aggrecanase protein of the invention, under the control of known regulatory sequences. The transformed host cells are cultured and the aggrecanase proteins recovered and purified from the culture medium. The purified proteins are substantially free from other proteins with which they are co-produced as well as from other contaminants.

10 Suitable cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO). The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening, product production and purification are known in the art. See, e.g., Gething and Sambrook, Nature, 293:620-625 (1981), or alternatively, Kaufman et al, Mol. Cell. Biol., 5(7):1750-1759 (1985) or Howley et al, 15 U.S. Patent 4,419,446. Another suitable mammalian cell line, which is described in the accompanying examples, is the monkey COS-1 cell line. The mammalian cell CV-1 may also be suitable.

Bacterial cells may also be suitable hosts. For example, the various strains of E. coli (e.g., HB101, MC1061) are well-known as host cells in the field of biotechnology. 20 Various strains of B. subtilis, Pseudomonas, other bacilli and the like may also be employed in this method. For expression of the protein in bacterial cells, DNA encoding the propeptide of Aggrecanase is generally not necessary.

Many strains of yeast cells known to those skilled in the art may also be available as host cells for expression of the polypeptides of the present invention. Additionally, where desired, insect cells may be utilized as host cells in the method of the present invention. See, e.g. Miller et al, Genetic Engineering, 8:277-298 (Plenum Press 1986) and references cited therein.

Another aspect of the present invention provides vectors for use in the method of expression of these novel aggrecanase polypeptides. Preferably the vectors contain the full novel DNA sequences described above which encode the novel factors of the invention. Additionally, the vectors contain appropriate expression control sequences permitting expression of the aggrecanase protein sequences. Alternatively, vectors incorporating modified sequences as described above are also embodiments of the present invention. Additionally, the sequence of SEQ ID NO:3 or SEQ ID NO: 1 or other sequences encoding aggrecanase proteins could be manipulated to express composite aggrecanase molecules. Thus, the present invention includes chimeric DNA molecules encoding an aggrecanase protein comprising a fragment from SEQ ID NO: 3 or SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO 1 linked in correct reading frame to a DNA sequence encoding another aggrecanase polypeptide.

The vectors may be employed in the method of transforming cell lines and contain selected regulatory sequences in operative association with the DNA coding sequences of the invention which are capable of directing the replication and expression thereof in selected host cells. Regulatory sequences for such vectors are known to those skilled in

the art and may be selected depending upon the host cells. Such selection is routine and does not form part of the present invention.

Various conditions such as osteoarthritis are known to be characterized by degradation of aggrecan. Therefore, an aggrecanase protein of the present invention
5 which cleaves aggrecan may be useful for the development of inhibitors of aggrecanase. The invention therefore provides compositions comprising an aggrecanase inhibitor. The inhibitors may be developed using the aggrecanase in screening assays involving a mixture of aggrecan substrate with the inhibitor followed by exposure to aggrecan. The compositions may be used in the treatment of osteoarthritis and other conditions
10 exhibiting degradation of aggrecan.

The invention further includes antibodies which can be used to detect aggrecanase and also may be used to inhibit the proteolytic activity of aggrecanase.

The therapeutic methods of the invention includes administering the aggrecanase inhibitor compositions topically, systemically, or locally as an implant or device. The
15 dosage regimen will be determined by the attending physician considering various factors which modify the action of the aggrecanase protein, the site of pathology, the severity of disease, the patient's age, sex, and diet, the severity of any inflammation, time of administration and other clinical factors. Generally, systemic or injectable administration will be initiated at a dose which is minimally effective, and the dose will be increased
20 over a preselected time course until a positive effect is observed. Subsequently, incremental increases in dosage will be made limiting such incremental increases to such levels that produce a corresponding increase in effect, while taking into account any

adverse affects that may appear. The addition of other known factors, to the final composition, may also effect the dosage.

Progress can be monitored by periodic assessment of disease progression. The progress can be monitored, for example, by x-rays, MRI or other imaging modalities, 5 synovial fluid analysis, and/or clinical examination.

The following examples illustrate practice of the present invention in isolating and characterizing human aggrecanase and other aggrecanase-related proteins, obtaining the human proteins and expressing the proteins via recombinant techniques.

EXAMPLES

10 **EXAMPLE 1**

Isolation of DNA

Potential novel aggrecanase family members were identified using a database screening approach. Aggrecanase-1 [Science284:1664-1666 (1999)] has at least six domains: signal, propeptide, catalytic domain, disintegrin, tsp and c-terminal. The 15 catalytic domain contains a zinc binding signature region, TAAHELGHVKF and a "MET turn" which are responsible for protease activity. Substitutions within the zinc binding region in the number of the positions still allow protease activity, but the histidine (H) and glutamic acid (E) residues must be present. The thrombospondin domain of Aggrecanase-1 is also a critical domain for substrate recognition and cleavage. It is these 20 two domains that determine our classification of a novel aggrecanase family member. The protein sequence of the Aggrecanase-1 DNA sequence was used to query against the

GeneBank ESTs focusing on human ESTs using TBLASTN. The resulting sequences were the starting point in the effort to identify full length sequence for potential family members. The nucleotide sequence of the aggrecanase of the present invention is comprised of one EST (AA588434) that contains homology over the catalytic domain and zinc binding motif of Aggrecanase-1(ADAMTS4).

This human aggrecanase sequence was isolated from a dT-primed cDNA library constructed in the plasmid vector pED6-dpc2. cDNA was made from human testes RNA purchased from Clontech. The probe to isolate the aggrecanase of the present invention was generated from the sequence obtained from the database search. The sequence of the probe was as follows: 5'-GGTCAAATCGCGTCAGTGTAATACGGG-3'. The DNA probe was radioactively labeled with ^{32}P and used to screen the human testes dT-primed cDNA library, under high stringency hybridization/washing conditions, to identify clones containing sequences of the human candidate #8.

Fifty thousand library transformants were plated at a density of approximately 5000 transformants per plate on 10 plates. Nitrocellulose replicas of the transformed colonies were hybridized to the ^{32}P labeled DNA probe in standard hybridization buffer (1X Blotto[25X Blotto = %5 nonfat dried milk, 0.02% azide in dH₂O] + 1% NP-40 + 6X SSC + 0.05% Pyrophosphate) under high stringency conditions (65°C for 2 hours shaking). After 2 hours hybridization, the radioactively labeled DNA probe containing hybridization solution was removed and the filters were washed under high stringency conditions (3X SSC, 0.05% Pyrophosphate for 5 minutes at RT; followed by 2.2X SSC, 0.05% Pyrophosphate for 15 minutes at RT; followed by 2.2X SSC, 0.05% Pyrophosphate for 1-2 minutes at 65°C). The filters were wrapped in Saran wrap and

exposed to X-ray film for overnight. The autoradiographs were developed and positively hybridizing transformants of various signal intensities were identified. These positive clones were picked; grown for 12 hours in selective medium(L-broth plus 100µg/ml ampicillin) and plated at low density (approximately 100 colonies per plate).

- 5 Nitrocellulose replicas of the colonies were hybridized to the ^{32}P labeled probe in standard hybridization buffer ((1X Blotto[25X Blotto = %5 nonfat dried milk, 0.02% azide in dH₂O] + 1% NP-40 + 6X SSC +0.05% Pyrophosphate) under high stringency conditions (65°C for 2 hours). After 2 hours hybridization, the radioactively labeled DNA probe containing hybridization solution was removed and the filters were washed
- 10 under high stringency conditions (3X SSC, 0.05% Pyrophosphate for 5 minutes at RT; followed by 2.2X SSC, 0.05% Pyrophosphate for 15 minutes at RT; followed by 2.2X SSC, 0.05% Pyrophosphate for 1-2 minutes at 65°C). The filters were wrapped in Saran wrap and exposed to X-ray film for overnight. The autoradiographs were developed and positively hybridizing transformants were identified. Bacterial stocks of purified
- 15 hybridization positive clones were made and plasmid DNA was isolated. The sequence of the cDNA insert was determined and is set forth in SEQ ID NO. 1 from nucleotide # 1086 (TCG) through # 3396 (CGC). This sequence has been deposited in the American Type Culture Collection 10801 University Blvd. Manassas, VA 20110-2209 USA as PTA -2284. The cDNA insert contained the sequences of the DNA probe used in the
- 20 hybridization. The 5'(prime) and 3' (prime) sequences of this isolated sequence was then extracted using the RACE protocol. The fully determined sequence is set forth in SEQ ID NO: 1 from nucleotide # 1 to # 3766 .

The human candidate #8 sequence obtained aligns with several ESTs in the public database. Candidate #8 shows homology with ADAMTS 7 and 6. The aggrecanase of the present invention contains the zinc binding signature region, a "MET turn", and tsp type -1 motif, however is missing the signal and propeptide regions and c-terminal spacer regions. It is with these criteria that candidate #8 is considered a novel Aggrecanase family member.

The aggrecanase sequence of the invention can be used to design probes for further screening for full length clones containing the isolated sequence.

The 5P (signal and propeptide) and 3P (C-terminal spacer regions) ends of the full-length version of EST8 were determined by RACE PCR using the Clontech Marathon cDNA Amplification Kit. The testes and stomach Marathon cDNA sources were used as substrates for the RACE reactions. 5P RACE primers used in the reactions were; GSP1 – AGTCTAGAAAGCTGGTGATGTAGTCACGGC and GSP2 – TAGATGCATATGTCATAGCGTGTGATGAGCACTGC (contains a Nsi1 site). The Advantage-2 PCR Kit from Clontech was used to set up nested RACE reactions following instructions in the user manual for the Marathon cDNA Amplification Kit; the amount the GSP primers used was 0.2 pmol/ul of each PCR primer/ul of reaction mix.

GSP1 primer was used for the first round of PCR and GSP2 primer was used for the nested reaction. Products from the nested RACE reactions were digested with Nsi1 (on the GSP2 primer) and Not1 (on the AP2 primer provided in the Clontech kit and used in the nested RACE PCR) and ligated into the CS2+ vector cut with Nsi1 and Not1. Ligated products were transformed into ElectroMAX DH10B cells from Life Technologies.

Cloned RACE products were plated at low density (approximately 300 colonies per plate). Nitrocellulose replicas of the transformed colonies were hybridized to a ³²P labeled DNA probe in standard hybridization buffer (1X Blotto [25X Blotto = 5% nonfat dried milk, 0.02% azide], 1% NP-40, 6X SSC, 0.05% pyrophosphate) under high stringency conditions (65°C for 2 hours shaking). Sequence at the 5' end of candidate 8-1 was used as a DNA probe: CTCGAGTCTGGGAAGCACCGTTAACATCC.

After 2 hours, the hybridization solution (hybridization buffer containing 1x10⁶ cpm ³²P labeled DNA probe) was removed and the filters were washed under high stringency conditions (3X SSC, 0.05% pyrophosphate for 5 minutes at RT standing; followed by 2.2X SSC, 0.05% pyrophosphate for 15 minutes shaking at RT; followed by 2.2X SSC, 0.05% pyrophosphate for 1-2 minutes shaking at 65°C). The filters were covered with Saran Wrap and exposed to X-ray film overnight. The autoradiographs were developed and positively hybridizing transformants of various signal intensities were identified. These positive clones were picked and then grown for 12 hours in selective medium (L-broth plus 100ug/ml ampicillin). Plasmid DNA was prepared and sent for DNA sequence analysis. A second round of hybridizations was performed using a probe that was made to sequence more 5' than candidate 8-1. The DNA probe sequence was deduced from the 5' RACE products. The second probe sequence was as follows: GAAGGCGATCTCATAGCTCTCCAGACT. Cloned RACE products were again plated and the same hybridization protocol was followed, except using the more 5' probe. The initiator Met was deduced from a consensus sequence derived from the 5' RACE products generated from both the testes and the stomach cDNAs. 3' RACE primers used

were; GSP1 – GCTCTAGACTGGTCTGAGTGCACCCCCAGCT and GSP2 –
GTCCTTTGCAAGAGCGCAGACCAC. The Advantage GC-2 PCR Kit from Clontech
was used to set up nested RACE reactions. Reactions were set up following the
instructions in the user manual for the Marathon cDNA Amplification Kit; with the
5 exception that the amount of GC melt used was 5ul per 50ul reaction; the amount the GSP
primers used was 0.2 pmol/ul of each PCR primer/ul of reaction mix. GSP1 primer was
used for the first round of PCR and GSP2 primer was used for the nested reaction.
Products from the nested RACE reactions were ligated into the pT-Adv vector using the
AdvanTAge PCR Cloning Kit, per manufacturer's instructions. Ligated products were
10 transformed into ElectroMAX DH10B cells from Life Technologies. Cloned RACE
products were plated at low density (approximately 300 colonies per plate).
Nitrocellulose replicas of the transformed colonies were hybridized to a ³²P labeled DNA
probe in standard hybridization buffer (1X Blotto [25X Blotto = 5% nonfat dried milk,
0.02% azide], 1% NP-40, 6X SSC, 0.05% pyrophosphate) under high stringency
15 conditions (65°C for 2 hours shaking). Sequence at the 3P end of candidate 8-1 was used
as a DNA probe: GCACTGTGCAGAGCACTCACCCCA. After 2 hours, the
hybridization solution (hybridization buffer containing 1x10⁶ cpm ³²P labeled DNA
probe) was removed and the filters were washed under high stringency conditions (3X
SSC, 0.05% pyrophosphate for 5 minutes at RT standing; followed by 2.2X SSC, 0.05%
20 pyrophosphate for 15 minutes shaking at RT; followed by 2.2X SSC, 0.05%
pyrophosphate for 1-2 minutes shaking at 65°C). The filters were covered with Saran
Wrap and exposed to X-ray film overnight. The autoradiographs were developed and

positively hybridizing transformants of various signal intensities were identified. The positive clones were picked and then grown for 12 hours in selective medium (L-broth plus 100ug/ml ampicillin). Plasmid DNA was prepared and sent for DNA sequence analysis. The stop codon was deduced from a consensus sequence derived from the 3P

5 RACE products generated from both the testes and the stomach cDNAs.

With the exception of the region from base pair 1332 to 1517 (for this description base pair #1 is A of the initiator Met (ATG), the full-length sequence of EST8 was confirmed. A search of the public databases revealed a partial sequence for EST8, termed ADAMTS10. We used the sequence from this partial clone to construct the

10 contiguous region of our EST8 (base pair 1332 to 1517) with synthetic oligonucleotides.

The full-length sequence for EST8 (SEQ ID NO:3) was the consensus sequence derived from the hybridization positive candidate 8-1, the publicly available sequences representing EST8, and the PCR products from the Clontech testes and stomach cDNAs. The final EST8 expression construct was assembled from 4 EST8 specific fragments.

15 The 5P portion of EST8, from base pair 1 – 1342, was PCR amplified from a pool of stomach and testes cDNAs and will be termed fragment 1. The following primers were used; 5P PCR primer -

AAATGGGCGAATTCCCACCATGGCTCCCGCCTGCCAGATCCTCCG (contains an 8 base pair linker (AAATGGGC) an EcoRI cloning site (GAATTC) and a Kozak

20 sequence (CCACC) upstream of the initiator Met) and 3P PCR primer -

CCGAGTCTAGAAAGCTGGTGATGTAG (contains an XbaI site (TCTAGA)). This PCR product was digested with EcoRI and XbaI using standard digestion conditions.

The next portion of the gene, fragment 2, was constructed using synthetic oligonucleotides. The synthetic fragment stretched from an Xba1 site to a BsrF1 site representing base pair 1333 to 1517 of EST8. The synthetic oligonucleotides consisted of the following sequence: the top strand consisted of –

- 5 CTAGACTCGGGCCTGGGGCTCTGCCTGAACAACCGACCCCCCAGACAGGACT
TTGTGTACCCGACAGTGGCACCGGGCCAAGCCTACGATGCAGATGAGCAATG
CCGCTTTCAGCATGGAGTCAAATCGCGTCAGTGTAATAACGGGGAGGTCTGC
AGCGAGCTGTGGTGTCTGAGCAAGAGCAA; the bottom strand consisted of –
CCGGTTGCTCTTGCTCAGACACCACAGCTCGCTGCAGACCTCCCCGTATTTAC
10 ACTGACGCGATTTGACTCCATGCTGAAAGCGGCATTGCTCATCTGCATCGTAG
GCTTGGCCCCGGTGCCACTGTCGGGTACACAAAGTCCTGTCTGGGGGGTTCGGTT
GTTCAAGCAGAGCCCCAGGCCCGAGT. The next portion of EST8, fragment 3, was
a BsrF1 to Sph1 fragment digested from candidate 8-1. This represented from base pair
1518 to 2783 of the full-length version of EST8. The 3P portion of EST8, termed
15 fragment 4 (base pair 2663 to 3314), was PCR amplified. The following primers were
used; 5P – GGGTTGTAGGGAAGTGGTCGCTCTG (located within fragment 3,
upstream of the Sph1 site) and 3P –
AAATGGGCCTCGAGCCCTAGTGGCCCTGGCAGGTTTTGC (contains an 8 base
pair linker (AAATGGGC) and a Xho1 site (CTCGAG) downstream of the stop codon
20 (TAG)). This PCR product was digested with Sph1 and Xho1 using standard digestion
conditions. A full-length version of EST8 was constructed by ligating these 4 described
fragments, 5P fragment 1 (EcoRI/Xba1), internal fragment 2 (Xba1/BsrF1), internal

fragment 3 (BsrF1/SphI), and 3P fragment 4 (SphI/XhoI) into the Cos expression vector pED6-dpc2 (digested with EcoRI and XhoI). The final construct had a mutation in the XhoI cloning site, which was destroyed in the ligation. This did not effect the EST8 coding sequence and was left in the construct.

5 **EXAMPLE 2**

Expression of Aggrecanase

In order to produce murine, human or other mammalian aggrecanase-related proteins, the DNA encoding it is transferred into an appropriate expression vector and introduced into mammalian cells or other preferred eukaryotic or prokaryotic hosts including insect host cell culture systems by conventional genetic engineering techniques. Expression system for biologically active recombinant human aggrecanase is contemplated to be stably transformed mammalian cells, insect, yeast or bacterial cells.

One skilled in the art can construct mammalian expression vectors by employing the sequence of SEQ ID NO: 3 or the sequence of SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of SEQ ID NO 1 or other DNA sequences encoding aggrecanase-related proteins or other modified sequences and known vectors, such as pCD [Okayama et al., *Mol. Cell Biol.*, 2:161-170 (1982)], pJL3, pJL4 [Gough et al., *EMBO J.*, 4:645-653 (1985)] and pMT2 CXM.

The mammalian expression vector pMT2 CXM is a derivative of p91023(b) (Wong et al., *Science* 228:810-815, 1985) differing from the latter in that it contains the ampicillin resistance gene in place of the tetracycline resistance gene and further contains

a XhoI site for insertion of cDNA clones. The functional elements of pMT2 CXM have been described (Kaufman, R.J., 1985, Proc. Natl. Acad. Sci. USA 82:689-693) and include the adenovirus VA genes, the SV40 origin of replication including the 72 bp enhancer, the adenovirus major late promoter including a 5' splice site and the majority of the adenovirus tripartite leader sequence present on adenovirus late mRNAs, a 3' splice acceptor site, a DHFR insert, the SV40 early polyadenylation site (SV40), and pBR322 sequences needed for propagation in E. coli.

Plasmid pMT2 CXM is obtained by EcoRI digestion of pMT2-VWF, which has been deposited with the American Type Culture Collection (ATCC), Rockville, MD (USA) under accession number ATCC 67122. EcoRI digestion excises the cDNA insert present in pMT2-VWF, yielding pMT2 in linear form which can be ligated and used to transform E. coli HB 101 or DH-5 to ampicillin resistance. Plasmid pMT2 DNA can be prepared by conventional methods. pMT2 CXM is then constructed using loopout/in mutagenesis [Morinaga, et al., Biotechnology 84: 636 (1984). This removes bases 1075 to 1145 relative to the Hind III site near the SV40 origin of replication and enhancer sequences of pMT2. In addition it inserts the following sequence:

5' PO-CATGGGCAGCTCGAG-3'

at nucleotide 1145. This sequence contains the recognition site for the restriction endonuclease Xho I. A derivative of pMT2CXM, termed pMT23, contains recognition sites for the restriction endonucleases PstI, Eco RI, SalI and XhoI. Plasmid pMT2 CXM and pMT23 DNA may be prepared by conventional methods.

pEMC2 β 1 derived from pMT21 may also be suitable in practice of the invention. pMT21 is derived from pMT2 which is derived from pMT2-VWF. As described above EcoRI digestion excises the cDNA insert present in pMT-VWF, yielding pMT2 in linear form which can be ligated and used to transform E. Coli HB 101 or DH-5 to ampicillin resistance. Plasmid pMT2 DNA can be prepared by conventional methods.

pMT21 is derived from pMT2 through the following two modifications. First, 76 bp of the 5' untranslated region of the DHFR cDNA including a stretch of 19 G residues from G/C tailing for cDNA cloning is deleted. In this process, a XhoI site is inserted to obtain the following sequence immediately upstream from DHFR: 5' -

10 CTGCAGGCGAGCCTGAATTCCTCGAGCCATCATG-3'

PstI

Eco RI XhoI

Second, a unique ClaI site is introduced by digestion with EcoRV and XbaI, treatment with Klenow fragment of DNA polymerase I, and ligation to a ClaI linker (CATCGATG). This deletes a 250 bp segment from the adenovirus associated RNA (VAI) region but does not interfere with VAI RNA gene expression or function. pMT21 is digested with EcoRI and XhoI, and used to derive the vector pEMC2B1.

A portion of the EMCV leader is obtained from pMT2-ECAT1 [S.K. Jung, et al, J. Virol 63:1651-1660 (1989)] by digestion with Eco RI and PstI, resulting in a 2752 bp fragment. This fragment is digested with TaqI yielding an Eco RI-TaqI fragment of 508 bp which is purified by electrophoresis on low melting agarose gel. A 68 bp adapter and

its complementary strand are synthesized with a 5' TaqI protruding end and a 3' XhoI protruding end which has the following sequence:

5'-CGAGGTTAAAAACGTCTAGGCCCCCGAACCACGGGGACGTGGTTTTCCTTT

TaqI

5

GAAAAACACGATTGC-3'

XhoI

This sequence matches the EMC virus leader sequence from nucleotide 763 to 827. It also changes the ATG at position 10 within the EMC virus leader to an ATT and is followed by a XhoI site. A three way ligation of the pMT21 Eco RI-16hoI fragment, the EMC virus EcoRI-Taql fragment, and the 68 bp oligonucleotide adapter TaqI-16hoI adapter resulting in the vector pEMC2 β 1.

This vector contains the SV40 origin of replication and enhancer, the adenovirus major late promoter, a cDNA copy of the majority of the adenovirus tripartite leader sequence, a small hybrid intervening sequence, an SV40 polyadenylation signal and the adenovirus VA I gene, DIIFR and β -lactamase markers and an EMC sequence, in appropriate relationships to direct the high level expression of the desired cDNA in mammalian cells.

The construction of vectors may involve modification of the aggrecanase-related DNA sequences. For instance, aggrecanase cDNA can be modified by removing the non-coding nucleotides on the 5' and 3' ends of the coding region. The deleted non-coding

nucleotides may or may not be replaced by other sequences known to be beneficial for expression. These vectors are transformed into appropriate host cells for expression of aggrecanase-related proteins. Additionally, the sequence of SEQ ID NO: 3 or SEQ ID NO: 1 comprising nucleotide # 1 to # 3766 or comprising nucleotide # 1086 to #3396 of
5 SEQ ID NO 1 or other sequences encoding aggrecanase-related proteins can be manipulated to express a mature aggrecanase-related protein by deleting aggrecanase encoding propeptide sequences and replacing them with sequences encoding the complete propeptides of other aggrecanase proteins.

One skilled in the art can manipulate the sequences of SEQ ID NO: 3 or SEQ ID
10 NO: 1 by eliminating or replacing the mammalian regulatory sequences flanking the coding sequence with bacterial sequences to create bacterial vectors for intracellular or extracellular expression by bacterial cells. For example, the coding sequences could be further manipulated (e.g. ligated to other known linkers or modified by deleting non-coding sequences therefrom or altering nucleotides therein by other known techniques).
15 The modified aggrecanase-related coding sequence could then be inserted into a known bacterial vector using procedures such as described in T. Taniguchi et al., Proc. Natl Acad. Sci. USA, 77:5230-5233 (1980). This exemplary bacterial vector could then be transformed into bacterial host cells and a aggrecanase-related protein expressed thereby. For a strategy for producing extracellular expression of aggrecanase-related proteins in
20 bacterial cells, see, e.g. European patent application EPA 177,343.

Similar manipulations can be performed for the construction of an insect vector [See, e.g. procedures described in published European patent application 155,476] for

expression in insect cells. A yeast vector could also be constructed employing yeast regulatory sequences for intracellular or extracellular expression of the factors of the present invention by yeast cells. [See, e.g., procedures described in published PCT application WO86/00639 and European patent application EPA 123,289].

5 A method for producing high levels of a aggrecanase-related protein of the invention in mammalian, bacterial, yeast or insect host cell systems may involve the construction of cells containing multiple copies of the heterologous Aggrecanase-related gene. The heterologous gene is linked to an amplifiable marker, e.g. the dihydrofolate reductase (DHFR) gene for which cells containing increased gene copies can be selected
10 for propagation in increasing concentrations of methotrexate (MTX) according to the procedures of Kaufman and Sharp, J. Mol. Biol., 159:601-629 (1982). This approach can be employed with a number of different cell types.

For example, a plasmid containing a DNA sequence for an aggrecanase-related protein of the invention in operative association with other plasmid sequences enabling
15 expression thereof and the DHFR expression plasmid pAdA26SV(A)3 [Kaufman and Sharp, Mol. Cell. Biol., 2:1304 (1982)] can be co-introduced into DHFR-deficient CHO cells, DUKX-BII, by various methods including calcium phosphate coprecipitation and transfection, electroporation or protoplast fusion. DHFR expressing transformants are selected for growth in alpha media with dialyzed fetal calf serum, and subsequently
20 selected for amplification by growth in increasing concentrations of MTX (e.g. sequential steps in 0.02, 0.2, 1.0 and 5uM MTX) as described in Kaufman et al., Mol Cell Biol., 5:1750 (1983). Transformants are cloned, and biologically active aggrecanase expression

is monitored by the assays described above. Aggrecanase protein expression should increase with increasing levels of MTX resistance. Aggrecanase polypeptides are characterized using standard techniques known in the art such as pulse labeling with [35S] methionine or cysteine and polyacrylamide gel electrophoresis. Similar procedures
5 can be followed to produce other related aggrecanase-related proteins.

In one example the aggrecanase gene of the present invention set forth in SEQ ID NO:3 is cloned into the expression vector pED6 [Kaufman et al., Nucleic Acid Res. 19:44885-4490(1991)]. COS and CHO DUKX B11 cells are transiently transfected with the aggrecanase sequence of the invention (+/- co-transfection of PACE on a separate
10 pED6 plaasmid) by lipofection(LF2000, Invitrogen). Duplicate tranfections are performed for each gene of interest: (a) one for harvesting conditioned media for activity assay and (b) one for 35-S-methionine/cysteine metabolic labeling.

On day one media is changed to DME(COS) or alpha(CHO) media + 1% heat-inactivated fetal calf serum+/- 100µg/ml heparin on wells(a) to be harvested for activity
15 assay. After 48h (day4), conditioned media is harvested for activity assay.

On day 3, the duplicate wells(b) were changed to MEM (methiooine-free/cysteine free) media + 1% heat-inactivated fetal callf serum +100µg/ml heparin + 100µCi/ml 35S-methiooine/cysteine (Redivue Pro mix, Amersham). Following 6h incubation at 37°C, conditioned media was harvested and run on SDS-PAGE gels under
20 reducing conditions. Proteins are visualized by autoradiography.

EXAMPLE 3**Biological Activity of Expressed Aggrecanase**

To measure the biological activity of the expressed aggrecanase-related proteins obtained in Example 2 above, the proteins are recovered from the cell culture and purified
5 by isolating the aggrecanase-related proteins from other proteinaceous materials with which they are co-produced as well as from other contaminants. The purified protein may be assayed in accordance with assays described above. Purification is carried out using standard techniques known to those skilled in the art.

Protein analysis is conducted using standard techniques such as SDS-PAGE
10 acrylamide [Laemmli, Nature 227:680 (1970)] stained with silver [Oakley, et al. Anal. Biochem. 105:361 (1980)] and by immunoblot [Towbin, et al. Proc. Natl. Acad. Sci. USA 76:4350 (1979)].

The foregoing descriptions detail presently preferred embodiments of the present invention. Numerous modifications and variations in practice thereof are expected to
15 occur to those skilled in the art upon consideration of these descriptions. Those modifications and variations are believed to be encompassed within the claims appended hereto.

What is claimed is:

1. An isolated DNA molecule comprising a DNA sequence set forth in SEQ ID NO. **3.**
2. An isolated DNA molecule comprising a DNA sequence set forth in SEQ ID NO. 1 from nucleotide #1086 to # 3396.
3. An isolated DNA molecule comprising a DNA sequence selected from the group consisting of
 - a) the sequence of SEQ ID NO. 3,
 - b) the sequence of SEQ ID NO: 1 comprising nucleotide # 1086 to #3396 naturally occurring human allelic sequences and equivalent degenerative codon sequences of a) and b).
4. A vector comprising a DNA molecule of claim 1 in operative association with an expression control sequence therefor.
5. A vector comprising a DNA molecule of claim 2 in operative association with an expression control sequence therefor.
6. A host cell transformed with the DNA sequence of claim 1.

7. A host cell transformed with a DNA sequence of claim 2.
8. A method for producing a purified human aggrecanase protein, said method comprising the steps of:
 - (a) culturing a host cell transformed with a DNA molecule according to claim 1;
 - and
 - (b) recovering and purifying said aggrecanase protein from the culture medium.
9. A method for producing a purified human aggrecanase protein, said method comprising the steps of:
 - (a) culturing a host cell transformed with a DNA molecule according to claim 2;
 - and
 - (b) recovering and purifying said aggrecanase protein from the culture medium.
10. A purified aggrecanase polypeptide comprising the amino acid sequence of SEQ ID NO:4.
11. A purified aggrecanase polypeptide comprising the amino acid sequence set forth in SEQ ID NO 2.
12. A purified aggrecanase polypeptide produced by the steps of

- (a) culturing a cell transformed with a DNA molecule according to claim 1;
and
 - (b) recovering and purifying from said culture medium a polypeptide
comprising the amino acid sequence set forth in SEQ ID NO.4.
13. A purified aggrecanase polypeptide produced by the steps of
- (a) culturing a cell transformed with a DNA molecule according to claim 2; and
 - (b) recovering and purifying from said culture medium a polypeptide
comprising the amino acid sequence set forth in SEQ ID NO. 2.
14. An antibody that binds to a purified aggrecanase protein of claim 10.
15. A method for developing inhibitors of aggrecanase comprising the use of
aggrecanase protein set forth in SEQ ID NO. 4 or a fragment thereof.
16. A method for developing inhibitors of aggrecanase comprising the use of
aggrecanase protein set forth in SEQ ID NO. 2 or a fragment thereof.
17. The method of claim 15 or 16 wherein said method comprises three dimensional
structural analysis.

18. The method of claim 15 or 16 wherein said method comprises computer aided drug design.
19. A composition for inhibiting the proteolytic activity of aggrecanase comprising a peptide molecule which binds to the aggrecanase inhibiting the proteolytic
5 degradation of aggrecan.
20. A method for inhibiting the cleavage of aggrecan in a mammal comprising administering to said mammal an effective amount of a compound that inhibits aggrecanase activity.

SEQUENCE LISTING

<110> Genetics Institute, Inc.

<120> NOVEL AGGRECANASE MOLECULES

<130> GI 5454

<150> 60/242,317

<151> 2000-10-20

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3766

<212> DNA

<213> unknown

<400> 1

```

gcgggcgctg aattctaggg agggccccggg cgcggcgag gctccaaaga agaagaaacc 60
aaggccccaga gaggagggc caggtgcagg gagcaggcga gggaaggatc cgtacagggg 120
cccaacacta ctccaccaac cgaagcccc aaaaggagcc cggatgatgt gcgaaggctg 180
tgaacagggg aggcggcact gtgggggctg ccggcagccg gggctgggga gagacatgtg 240
gacacgtggc ctctatggct cccgcctgcc agatcctcgg ctgggcccctc gccctggggc 300
tgggcctcat gttcgaggtc acgcacgcct tccggtctca agatgagttc ctgtccagtc 360
tgagagagcta tgagatgcgc tccccaccc gcgtggacca caacggggca ctgctggcct 420
tctgcacacc tctccccgg aggcagcgcc gcggcagggg ggccacagcc gagtccccgc 480
tcttctacaa agtggcctcg cccagcaccc acttccctgt gaacctgacc cgcagctccc 540
gtctactggc agggcacgtc tccgtggagt actggacacg ggagggcctg gccctggcga 600
ggcgggcccc gcccactgc ctctacgtgt gtcacctgca gggccaggcc agcagctccc 660
atgtggccat cagcacctgt ggaggcctgc acggcctgat cgtggcagac gaggaagagt 720
acctgattga gccctgcac ggtgggcccc aggggttctc gagcccgag gaaagtggac 780
cacatgtggt gtacaagcgt tctctctctg tcacccccca cctggacaca gccgtggag 840
tgagagatga gaaacgtgg aaaggcgcc catggtggct gcggaccttg aagccaccgc 900
ctgcacggcc cctggggaat gaaacagagc gtggccagcc aggcctgaag cgatcgggtc 960
gccgagagcg ctacgtggag accctggttg tggctgacaa gatgatggtg gcctatcacg 1020
ggcgccggga tgtggagcag tatgtcctgg catcatgaa caltggtgcc aaacttttcc 1080
aggactcgag tctgggaagc accgttaaca tctctgtaac tcgctcctc ctgtctcagg 1140
aggaccagcc cactctggag atcacccacc atgcggggaa gtccctggac agcttctgta 1200
agtggcagaa atccatcgtg aaccacagcg gccatggcaa tgccattcca gagaacgggtg 1260
tggctaacca tgacacagca gtgctcatca cagctatga catctgcac tacaagaaca 1320
aaccctgcgg cacactaggg ctggccccgg tgggcggaat gtgtgagcgc gagagaagct 1380
gcagcgtcaa tgaggacatt ggctgggcca cagcgttcac cattgcccac gagatcgggc 1440
acacattcgg catgaacct gacggcggtg gaaacagctg tggggcccgt ggtcaggacc 1500
cagccaagct catggctgcc cacattacca ccaattcgtg cccattcgtg tggctcctc 1560
gcagcgtga ctacatcacc agctttctag actcagggcc tggggctctg cctgaacaac 1620
cggcccccca gacaggactt tgtgtaccgg acagtggcac cggggccaagc ctacgatgca 1680
gatgagcaat gccgctttca gcatggagtc aaatcgcgtc agtgtaaata cgggaggtct 1740
gcagcgagct glggtgtctg agcaagagca accggtgcat caccaacagc atcccgccg 1800
ccgagggcac gctgtgccag acgcacacca tcgacaaggg gtggtgctac aaacgggtct 1860
gtgtccccct tgggtgcgc ccagaggggtg tggacggagc ctggggggcg tggactccat 1920
ggggcgactg cagccggacc tgtggcgggc gcgtgtctc ttctagccgt cactgcgaca 1980
gccccaggcc aaccatcggt ggcaagtact gtctgggtga gagaaggcgg caccgctcct 2040
gcaacacgga tgactgtccc cctggctccc aggaacttcag agaagtgcag tgttctgaat 2100
ttgacagcat ccttttccgt gggaaattct acaagtggaa aacgtaccgg ggagggggcg 2160
tgaaggcctg ctgcctcacc tgccatagcgg aaggttcaa cttctacacg gagagggcg 2220
cagcgtgtgt cagcgggaca cctgcctgac cagacacggt ggacatttgc gtcagtggcg 2280
aatgcaagca cgtgggctgc gaccgagtc tgggctccga cctgcgggag gacaagtgc 2340
caatgtgtg caatgacac aatgcctacg agaccatcga ggcgctctc agcccagcct 2400

```

```

atgcccccat cccccgtgac tgcctgcccc tctactcctg caactatcgg ccttggacca 2460
aqtgctcggc caagtggtga ggggtgagcc aqgtgcaggg ggtggagtgc cggacccagg 2520

```

```

tggacagetc cgcggtegec cccactact gcagtgccca cagcaagctg cccaaaaggc 2880
agcgcgcctg caacacggag ccttgccctc cagactgggt ttaggggaac tggtcgctct 2940
gcagccgcag ctgcgatgca ggcgtgcgca gccgctcggt cgtgtgccag cgcgcgctct 3000
ctgcgcgga ggagaaggcg ctggacgaca gcgcatgccc gcagccgcgc ccacctgtac 3060
tggaggcctg ccacggcccc acttgccctc cggagtgggc ggccctcgac tggctcgagt 3120
gcacccccag ctgcggggcg ggccctcgcc accgcgtggc cctttgcaag agcgcagacc 3180
accgcgccac gctgcccccg gcgcaactgt caccgcgcgc caagccaccg gccaccatgc 3240
gctgcaactt gcgcgcgtgc ccccgggccc gctgggtggc tggcgagtgg ggtgagtgct 3300
ctgcacagtg cggcgteggg cagcggcagc gctcggtgcg ctgcaccagc cacacggggc 3360
aggcgtegca cgaagtgcag gaggcctgc ggccgcccac cacgcagcaa tgtgaggcca 3420
agtgcgacag cccaaccccc gggggcgggc ctgaagagtg caaggatgtg aacaaggctc 3480
cctactgccc cctggtgtgc aaatttcagt tctgcagccg agcctacttc cgccagatgt 3540
gtgcacaaac ctgccagggc cactaggggg cgcgcggcac ccggagccac agctggcggg 3600
gtctccgccc ccagtcctgc agcggggcgg ccagaggggg ccccgggggg gcgggaactg 3660
ggaggggaagg gtgagacgga gccggaagtt atttattggg aacctctgca gggccctggc 3720
tgggggggatg gagaggggct ggtatccac ctgcccgggc ggccgc 3766

```

<210> 2
 <211> 770
 <212> PRT
 <213> unknown

<220>
 <221> VARIANT
 <222> (1)...(770)
 <223> Xaa = Any Amino Acid

<400> 2

Ser	Ser	Leu	Gly	Ser	Thr	Val	Asn	Ile	Leu	Val	Thr	Arg	Leu	Ile	Leu
1				5					10					15	
Leu	Thr	Glu	Asp	Gln	Pro	Thr	Leu	Glu	Ile	Thr	His	His	Ala	Gly	Lys
		20						25					30		
Ser	Leu	Asp	Ser	Phe	Cys	Lys	Trp	Gln	Lys	Ser	Ile	Val	Asn	His	Ser
		35					40					45			
Gly	His	Gly	Asn	Ala	Ile	Pro	Glu	Asn	Gly	Val	Ala	Asn	His	Asp	Thr
	50				55					60					
Ala	Val	Leu	Ile	Thr	Arg	Tyr	Asp	Ile	Cys	Ile	Tyr	Lys	Asn	Lys	Pro
	65				70				75						80
Cys	Gly	Thr	Leu	Gly	Leu	Ala	Pro	Val	Gly	Gly	Met	Cys	Glu	Arg	Glu
			85					90					95		
Arg	Ser	Cys	Ser	Val	Asn	Glu	Asp	Ile	Gly	Leu	Ala	Thr	Ala	Phe	Thr
			100					105					110		
Ile	Ala	His	Glu	Ile	Gly	His	Thr	Phe	Gly	Met	Asn	His	Asp	Gly	Val
		115					120					125			
Gly	Asn	Ser	Cys	Gly	Ala	Arg	Gly	Gln	Asp	Pro	Ala	Lys	Leu	Met	Ala
	130					135					140				
Ala	His	Ile	Thr	Met	Lys	Thr	Asn	Pro	Phe	Val	Trp	Ser	Ser	Cys	Ser
	145				150					155					160
Arg	Asp	Tyr	Ile	Thr	Ser	Phe	Leu	Asp	Ser	Gly	Pro	Gly	Ala	Leu	Pro
			165					170						175	
Glu	Gln	Pro	Ala	Pro	Gln	Thr	Gly	Leu	Cys	Val	Pro	Asp	Ser	Gly	Thr
		180						185					190		
Gly	Pro	Ser	Leu	Arg	Cys	Arg	Xaa	Ala	Met	Pro	Leu	Ser	Ala	Trp	Ser
		195					200					205			
Gln	Ile	Ala	Ser	Val	Xaa	Ile	Arg	Glu	Val	Cys	Ser	Glu	Leu	Trp	Cys
	210				215						220				
Leu	Ser	Lys	Ser	Asn	Arg	Cys	Ile	Thr	Asn	Ser	Ile	Pro	Ala	Ala	Glu
				230						235					240
Gly	Thr	Leu	Cys	Gln	Thr	His	Thr	Ile	Asp	Lys	Gly	Trp	Cys	Tyr	Lys
				245					250					255	
Arg	Val	Cys	Val	Pro	Phe	Gly	Ser	Arg	Pro	Glu	Gly	Val	Asp	Gly	Ala
			260					265					270		
Trp	Gly	Pro	Trp	Thr	Pro	Trp	Gly	Asp	Cys	Ser	Arg	Thr	Cys	Gly	Gly

Gly Gly Lys Tyr Cys Leu Gly Glu Arg Arg Arg His Arg Ser Cys Asn
 305 310 315 320
 Thr Asp Asp Cys Pro Pro Gly Ser Gln Asp Phe Arg Glu Val Gln Cys
 325 330 335
 Ser Glu Phe Asp Ser Ile Pro Phe Arg Gly Lys Phe Tyr Lys Trp Lys
 340 345 350
 Thr Tyr Arg Gly Gly Gly Val Lys Ala Cys Ser Leu Thr Cys Leu Ala
 355 360 365
 Glu Gly Phe Asn Phe Tyr Thr Glu Arg Ala Ala Val Val Asp Gly
 370 375 380
 Thr Pro Cys Arg Pro Asp Thr Val Asp Ile Cys Val Ser Gly Glu Cys
 385 390 395 400
 Lys His Val Gly Cys Asp Arg Val Leu Gly Ser Asp Leu Arg Glu Asp
 405 410 415
 Lys Cys Arg Val Cys Gly Gly Asp Gly Ser Ala Cys Glu Thr Ile Glu
 420 425 430
 Gly Val Phe Ser Pro Ala Ser Pro Gly Ala Gly Tyr Glu Asp Val Val
 435 440 445
 Trp Ile Pro Lys Gly Ser Val His Ile Phe Ile Gln Asp Leu Asn Leu
 450 455 460
 Ser Leu Ser His Leu Ala Leu Lys Gly Asp Gln Glu Ser Leu Leu Leu
 465 470 475 480
 Glu Gly Leu Pro Gly Thr Pro Gln Pro His Arg Leu Pro Leu Ala Gly
 485 490 495
 Thr Thr Phe Gln Leu Arg Gln Gly Pro Asp Gln Val Gln Ser Leu Glu
 500 505 510
 Ala Leu Gly Pro Ile Asn Ala Ser Leu Ile Val Met Val Leu Ala Arg
 515 520 525
 Thr Glu Leu Pro Ala Leu Arg Tyr Arg Phe Asn Ala Pro Ile Ala Arg
 530 535 540
 Asp Ser Leu Pro Pro Tyr Ser Trp His Tyr Ala Pro Trp Thr Lys Cys
 545 550 555 560
 Ser Ala Gln Cys Ala Gly Gly Ser Gln Val Gln Ala Val Glu Cys Arg
 565 570 575
 Asn Gln Leu Asp Ser Ser Ala Val Ala Pro His Tyr Cys Ser Ala His
 580 585 590
 Ser Lys Leu Pro Lys Arg Gln Arg Ala Cys Asn Thr Glu Pro Cys Pro
 595 600 605
 Pro Asp Trp Val Val Gly Asn Trp Ser Leu Cys Ser Arg Ser Cys Asp
 610 615 620
 Ala Gly Val Arg Ser Arg Ser Val Val Cys Gln Arg Arg Val Ser Ala
 625 630 635 640
 Ala Glu Glu Lys Ala Leu Asp Asp Ser Ala Cys Pro Gln Pro Arg Pro
 645 650 655
 Pro Val Leu Glu Ala Cys His Gly Pro Thr Cys Pro Pro Glu Trp Ala
 660 665 670
 Ala Leu Asp Trp Ser Glu Cys Thr Pro Ser Cys Gly Pro Gly Leu Arg
 675 680 685
 His Arg Val Val Leu Cys Lys Ser Ala Asp His Arg Ala Thr Leu Pro
 690 695 700
 Pro Ala His Cys Ser Pro Ala Ala Lys Pro Pro Ala Thr Met Arg Cys
 705 710 715 720
 Asn Leu Arg Arg Cys Pro Pro Ala Arg Trp Val Ala Gly Glu Trp Gly
 725 730 735
 Glu Cys Ser Ala Gln Cys Gly Val Gly Gln Arg Gln Arg Ser Val Arg
 740 745 750
 Cys Thr Ser His Thr Gly Gln Ala Ser His Glu Cys Thr Glu Ala Leu
 755 760 765
 Arg Pro
 770

<210> 3
 <211> 3377
 <212> DNA
 <213> Unknown

<220>
 <223> unknown

<400> 3
 gaattccac catggtccccc gcttgcacaga tctctccgtg ggcctctgcc ctggggcttg 60
 gctcatgtt cgagggtcac cagcctctcc ggtctcaaga tgagttccctg tccagttctg 120
 agagctatga gatcgccttc cccacccgcg tggaccacaa cggggcactg ctggccttct 180
 cgccaccttc tccccggagg cagcgccgcg gcacgggggc cacagccgag tcccgcctct 240
 tctacaaagt ggcctcgccc agcacccact tctgtctgaa cctgaccgcg agctcccgct 300
 tactggcagg gcacgtcttc gtggagtact ggacacggga gggcctggcc tggcagagag 360
 cggcccgccc cactgccttc tacgttggtc acctgcaggg ccaggccagc agctcccatg 420
 tggccatcag cactgttggg ggcctgcacg gcctgacgt ggacagcag gaagagtacc 480
 tgattgagcc cctgcacggg gggcccaagg gttctcggag cccggaggaa agtggaccac 540
 atgtggtgta caagcgttcc tctctgcgtc acccccacct ggacacagcc tgtggagtga 600
 gagatgagaa accgtggaaa gggcggccat ggtggctgcg gaccttgaag ccaccgcctg 660
 ccaggccctt ggggaatgaa acagagcgtg gcacgccagg cctgaagcga tcggtcagcc 720
 gagagcgcta ctgggagacc ctggtggtgg ctgacaagat gatggtggcc tatcacgggc 780
 gccgggaigt yyagcagtat gtcttggcca tcatgaacat tgttgccaaa cttttccagg 840
 actcgagttt ggggaagcacc gttaacatcc tctgaactcg cctcatcctg ctccagggag 900
 accagcccac tctggagatc acccaccatg ccgggaagtc cctggacagc tctgttaagt 960
 ggacagaatc catcgtgaac cacagcggtc atggcaatgc catlccagag aacggtgttg 1020
 ctaacctga cacagcagtg ctcatcacac gctatgacat ctgcatctac aagaacaaac 1080
 cctgcccgcac actaggcctg gccccggttg gcggaatgtg tgagcgcgag agaagctgca 1140
 cgttcaatga ggacattggc ctggccacag cgttcaccat tgcccacgag atcgggcaca 1200
 gattcggcat gaacctgac ggctggggaa agcagctgtg gggccgtggt caggaccag 1260
 ccaagctcat ggctgcccac attaccatga agaccaaccc gtctgtgttg tcatcctgca 1320
 gccgtgacta catcaccagc tttctagact cgggcctggg gctctgctg aacaaccgac 1380
 cccccagaca ggactttgtg taccgacag tggcaccggg ccaagcctac gatgcagatg 1440
 agcaatgccg ctttcagcat ggaagtcaat aagagcaacc ggtgcacac taaatacggg gaggtctgca 1500
 gcgagctgtg gtgtctgagc aagagcaacc ggtgcacac caacagcatc ccggcccgcc 1560
 agggcacgct gtgccagacg cacaccatcg acaaggggtg gtgctacaaa cgggtctgtg 1620
 tcccccttgg gtcgcgccca gagggtgttg acggagcctg ggggcccgtg actccatggg 1680
 gcgactgcag ccggacgtgt ggcggcgccg tctctcttc tagccgtcac tgcgacagcc 1740
 ccaggccaac catcgggggc aagtaactgt tgggtgagag aaggcgccac cgtcctgca 1800
 acacggatga ctgtcccccct ggctcccagg acttcagaga agtgacgtgt tctgaatttg 1860
 acagcatccc tttccgtggg aaattctaca agtggaatac gtaccgggga gggggcgtga 1920
 aggcactgct ctttcagctg ctacgggaag ctacacggag agggcgccag 1980
 ccgtggtgga cgggacaccc tgccgtccag acacggtgga catttgcgtc agtggcgaat 2040
 gcaagcacgt gggctgcgac cgaagccttg gctccgacct cggggaggac aagtgccag 2100
 tgtgtggcgg tgacggcagt gctgtctgga ccacagagg cgtcttcagc ccagcctcac 2160
 ctggggcccg gtacgaggat gtcgtctgga ttcccaagg ctcgctccac atcttcatcc 2220
 aggatctgaa cctctctctc agtcaacttg ccttgaagg agaccaggag tccctgctgc 2280
 tggaggggct gcccgggacc cccagcccc accgtctgcc tctagctggg accaccttcc 2340
 aactggcaca ggggccagac caggtccaga gctcgaagc cctgggaccg attaatgcat 2400
 ctctcatcgt catggtgctg gcccggaacc gctcgcctg cctccgctac cgcttcaatg 2460
 cccccatcgc ccgtgactcg ctgccccctt actcctggca ctatgcgccc tggaccaagt 2520
 gctcggccca gtgtgcaggg ggtagccagg tgcaaggcgt ggagtgccgc aaccagctgg 2580
 acagctccgc ggtcgcctcc cactactgca gtgccacag caagctgccc aaaaggcagc 2640
 ggcctgcaa cacggagcct tgccctccag actgggttgt agggaaactg tcgctctgca 2700
 gccgcagctg cgtatcaggg gtgcgcagcc gctcggctgt gtgccagcgc cgcgtctctg 2760
 ccgcggagga gaaggcgtg gacgacagcg catgcccgca gccgcgcccc cctgtactgg 2820
 aggctgccca cggcccaact tgccctccgg agtggggcgc cctcgactgg tctgagtga 2880
 cccccagttg ccggccgggg ctccgccacc gcgtggtcct ttgcaagagc gcagaccacc 2940
 gcgccacgtt gcccccgggc cactgtctac ccgccgcaa gccaccggcc accatgcgct 3000
 gcaacttgcg ccgtgcccc ccggcccgtt ggggtggtgg cgagtggggt gagtgtctct 3060
 cacagtgcgg cgtcggggcag cggcagcgtt cgggtgcgtg caccagccac acgggcccag 3120
 cgtcgcacga gtgcacggag gccctgcggc cggccaccac gcagcagtg gaggccaagt 3180
 gggccagccc gggcccgagg gggccgctg ggggtgcaa ggggtgcctt aaggtgcctt 3240

<210> 4
 <211> 1104
 <212> PRT
 <213> unknown

<400> 4

Met	Ala	Pro	Ala	Cys	Gln	Ile	Leu	Arg	Trp	Ala	Leu	Ala	Leu	Gly	Leu
1				5					10					15	
Gly	Leu	Met	Phe	Glu	Val	Thr	His	Ala	Phe	Arg	Ser	Gln	Asp	Glu	Phe
			20					25					30		
Leu	Ser	Ser	Leu	Glu	Ser	Tyr	Glu	Ile	Ala	Phe	Pro	Thr	Arg	Val	Asp
		35					40					45			
His	Asn	Gly	Ala	Leu	Leu	Ala	Phe	Ser	Pro	Pro	Pro	Pro	Arg	Arg	Gln
	50					55					60				
Arg	Arg	Gly	Thr	Gly	Ala	Thr	Ala	Glu	Ser	Arg	Leu	Phe	Tyr	Lys	Val
65					70					75					80
Ala	Ser	Pro	Ser	Thr	His	Phe	Leu	Leu	Asn	Leu	Thr	Arg	Ser	Ser	Arg
				85					90					95	
Leu	Leu	Ala	Gly	His	Val	Ser	Val	Glu	Tyr	Trp	Thr	Arg	Glu	Gly	Leu
			100					105					110		
Ala	Trp	Gln	Arg	Ala	Ala	Arg	Pro	His	Cys	Leu	Tyr	Ala	Gly	His	Leu
		115					120					125			
Gln	Gly	Gln	Ala	Ser	Ser	Ser	His	Val	Ala	Ile	Ser	Thr	Cys	Cly	Cly
	130					135					140				
Leu	His	Gly	Leu	Ile	Val	Ala	Asp	Glu	Glu	Glu	Tyr	Leu	Ile	Glu	Pro
145					150					155					160
Leu	His	Gly	Gly	Pro	Lys	Gly	Ser	Arg	Ser	Pro	Glu	Glu	Ser	Gly	Pro
				165					170					175	
His	Val	Val	Tyr	Lys	Arg	Ser	Ser	Leu	Arg	His	Pro	His	Leu	Asp	Thr
			180					185					190		
Ala	Cys	Gly	Val	Arg	Asp	Glu	Lys	Pro	Trp	Lys	Gly	Arg	Pro	Trp	Trp
	195						200					205			
Leu	Arg	Thr	Leu	Lys	Pro	Pro	Pro	Ala	Arg	Pro	Leu	Gly	Asn	Glu	Thr
	210					215					220				
Glu	Arg	Gly	Gln	Pro	Gly	Leu	Lys	Arg	Ser	Val	Ser	Arg	Glu	Arg	Tyr
225					230					235					240
Val	Glu	Thr	Leu	Val	Val	Ala	Asp	Lys	Met	Met	Val	Ala	Tyr	His	Gly
				245					250					255	
Arg	Arg	Asp	Val	Glu	Gln	Tyr	Val	Leu	Ala	Ile	Met	Asn	Ile	Val	Ala
			260					265					270		
Lys	Leu	Phe	Gln	Asp	Ser	Ser	Leu	Gly	Ser	Thr	Val	Asn	Ile	Leu	Val
	275						280					285			
Thr	Arg	Leu	Ile	Leu	Leu	Thr	Glu	Asp	Gln	Pro	Thr	Leu	Glu	Ile	Thr
	290					295					300				
His	His	Ala	Gly	Lys	Ser	Leu	Asp	Ser	Phe	Cys	Lys	Trp	Gln	Lys	Ser
305					310					315					320
Ile	Val	Asn	His	Ser	Gly	His	Gly	Asn	Ala	Ile	Pro	Glu	Asn	Gly	Val
				325					330					335	
Ala	Asn	His	Asp	Thr	Ala	Val	Leu	Ile	Thr	Arg	Tyr	Asp	Ile	Cys	Ile
			340					345					350		
Tyr	Lys	Asn	Lys	Pro	Cys	Gly	Thr	Leu	Gly	Leu	Ala	Pro	Val	Gly	Gly
		355					360					365			
Met	Cys	Glu	Arg	Glu	Arg	Ser	Cys	Ser	Val	Asn	Glu	Asp	Ile	Gly	Leu
	370					375					380				
Ala	Thr	Ala	Phe	Thr	Ile	Ala	His	Glu	Ile	Gly	His	Thr	Phe	Gly	Met
385					390					395					400
Asn	His	Asp	Gly	Val	Gly	Asn	Ser	Cys	Gly	Ala	Arg	Gly	Gln	Asp	Pro
				405					410					415	
Ala	Lys	Leu	Met	Ala	Ala	His	Ile	Thr	Met	Lys	Thr	Asn	Pro	Phe	Val
			420					425					430		
Trp	Ser	Ser	Cys	Ser	Arg	Asp	Tyr	Ile	Thr	Ser	Phe	Leu	Asp	Ser	Gly
		435					440					445			

Phe	Gln	His	Gly	Val	Lys	Ser	Arg	Gln	Cys	Lys	Tyr	Gly	Glu	Val	Cys
				485					490					495	
Ser	Glu	Leu	Trp	Cys	Leu	Ser	Lys	Ser	Asn	Arg	Cys	Ile	Thr	Asn	Ser
			500					505					510		
Ile	Pro	Ala	Ala	Glu	Gly	Thr	Leu	Cys	Gln	Thr	His	Thr	Ile	Asp	Lys
		515					520					525			
Gly	Trp	Cys	Tyr	Lys	Arg	Val	Cys	Val	Pro	Phe	Gly	Ser	Arg	Pro	Glu
	530					535					540				
Gly	Val	Asp	Gly	Ala	Trp	Gly	Pro	Trp	Thr	Pro	Trp	Gly	Asp	Cys	Ser
545					550					555					560
Arg	Thr	Cys	Gly	Gly	Gly	Val	Ser	Ser	Ser	Ser	Arg	His	Cys	Asp	Ser
				565					570					575	
Pro	Arg	Pro	Thr	Ile	Gly	Gly	Lys	Tyr	Cys	Leu	Gly	Glu	Arg	Arg	Arg
			580					585					590		
His	Arg	Ser	Cys	Asn	Thr	Asp	Asp	Cys	Pro	Pro	Gly	Ser	Gln	Asp	Phe
		595					600					605			
Arg	Glu	Val	Gln	Cys	Ser	Glu	Phe	Asp	Ser	Ile	Pro	Phe	Arg	Gly	Lys
	610					615					620				
Phe	Tyr	Lys	Trp	Lys	Thr	Tyr	Arg	Gly	Gly	Gly	Val	Lys	Ala	Cys	Ser
625					630					635					640
Leu	Thr	Cys	Leu	Ala	Glu	Gly	Phe	Asn	Phe	Tyr	Thr	Glu	Arg	Ala	Ala
				645					650					655	
Ala	Val	Val	Asp	Gly	Thr	Pro	Cys	Arg	Pro	Asp	Thr	Val	Asp	Ile	Cys
			660					665				670			
Val	Ser	Gly	Glu	Cys	Lys	His	Val	Gly	Cys	Asp	Arg	Val	Leu	Gly	Ser
		675					680					685			
Asp	Leu	Arg	Glu	Asp	Lys	Cys	Arg	Val	Cys	Gly	Gly	Asp	Gly	Ser	Ala
	690					695					700				
Cys	Glu	Thr	Ile	Glu	Gly	Val	Phe	Ser	Pro	Ala	Ser	Pro	Gly	Ala	Gly
705					710					715					720
Tyr	Glu	Asp	Val	Val	Trp	Ile	Pro	Lys	Gly	Ser	Val	His	Ile	Phe	Ile
				725					730					735	
Gln	Asp	Leu	Asn	Leu	Ser	Leu	Ser	His	Leu	Ala	Leu	Lys	Gly	Asp	Gln
			740					745					750		
Glu	Ser	Leu	Leu	Leu	Glu	Gly	Leu	Pro	Gly	Thr	Pro	Gln	Pro	His	Arg
		755					760					765			
Leu	Pro	Leu	Ala	Gly	Thr	Thr	Phe	Gln	Leu	Arg	Gln	Gly	Pro	Asp	Gln
	770					775					780				
Val	Gln	Ser	Leu	Glu	Ala	Leu	Gly	Pro	Ile	Asn	Ala	Ser	Leu	Ile	Val
785					790					795					800
Met	Val	Leu	Ala	Arg	Thr	Glu	Leu	Pro	Ala	Leu	Arg	Tyr	Arg	Phe	Asn
				805					810					815	
Ala	Pro	Ile	Ala	Arg	Asp	Ser	Leu	Pro	Pro	Tyr	Ser	Trp	His	Tyr	Ala
			820					825					830		
Pro	Trp	Thr	Lys	Cys	Ser	Ala	Gln	Cys	Ala	Gly	Gly	Ser	Gln	Val	Gln
		835					840					845			
Ala	Val	Glu	Cys	Arg	Asn	Gln	Leu	Asp	Ser	Ser	Ala	Val	Ala	Pro	His
	850					855					860				
Tyr	Cys	Ser	Ala	His	Ser	Lys	Leu	Pro	Lys	Arg	Gln	Arg	Ala	Cys	Asn
865					870					875					880
Thr	Glu	Pro	Cys	Pro	Pro	Asp	Trp	Val	Val	Gly	Asn	Trp	Ser	Leu	Cys
				885					890					895	
Ser	Arg	Ser	Cys	Asp	Ala	Gly	Val	Arg	Ser	Arg	Ser	Val	Val	Cys	Gln
			900					905					910		
Arg	Arg	Val	Ser	Ala	Ala	Glu	Glu	Lys	Ala	Leu	Asp	Asp	Ser	Ala	Cys
		915					920					925			
Pro	Gln	Pro	Arg	Pro	Pro	Val	Leu	Glu	Ala	Cys	His	Gly	Pro	Thr	Cys
	930					935						940			
Pro	Pro	Glu	Trp	Ala	Ala	Leu	Asp	Trp	Ser	Glu	Cys	Thr	Pro	Ser	Cys
945					950					955					960
Gly	Pro	Gly	Leu	Arg	His	Arg	Val	Val	Leu	Cys	Lys	Ser	Ala	Asp	His
				965					970					975	

1010	1015	1020
Gln Arg Ser Val Arg Cys Thr Ser His Thr Gly Gln Ala Ser His Glu		
1025	1030	1035
Cys Thr Glu Ala Leu Arg Pro Pro Thr Thr Gln Gln Cys Glu Ala Lys		
	1045	1050
Cys Asp Ser Pro Thr Pro Gly Asp Gly Pro Glu Glu Cys Lys Asp Val		
	1060	1065
Asn Lys Val Ala Tyr Cys Pro Leu Val Leu Lys Phe Gln Phe Cys Ser		
	1075	1080
Arg Ala Tyr Phe Arg Gln Met Cys Cys Lys Thr Cys Gln Gly His Xaa		
1090	1095	1100